

University of Dundee

Increased filaggrin expression in oral lichenoid lesions

Brown, S. J.

Published in:

Journal of the European Academy of Dermatology and Venereology

DOI:

[10.1111/jdv.14265](https://doi.org/10.1111/jdv.14265)

Publication date:

2017

Document Version

Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Brown, S. J. (2017). Increased filaggrin expression in oral lichenoid lesions: is this cause or effect? *Journal of the European Academy of Dermatology and Venereology*, 31(5), 759-759. <https://doi.org/10.1111/jdv.14265>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

This is the peer reviewed version of the following article: Brown, S.J. (2017), Increased filaggrin expression in oral lichenoid lesions: is this cause or effect?. *J Eur Acad Dermatol Venereol*, 31(5): 759. doi:10.1111/jdv.14265, which has been published in final form at doi:10.1111/jdv.14265. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving

JEADV Commentary

Increased filaggrin expression in oral lichenoid lesions: is this cause or effect?

Sara J Brown

Professor of Molecular & Genetic Dermatology, Wellcome Trust Senior Research Fellow in Clinical Science, Honorary Consultant Dermatologist

Skin Research Group,
School of Medicine,
Ninewells Hospital & Medical School,
Jacqui Wood Centre level 7,
James Arrott Drive,
University of Dundee,
Dundee DD1 9SY

T +44 (0) 1382 383210

E s.j.brown@dundee.ac.uk

Filaggrin is expressed in the outermost layers of the human epidermis;¹ it plays a central role in skin barrier formation through its multiple functions from the pro-protein which has calcium-binding domains, to the filaggrin monomers which aggregate keratin filaments, and to its constituent amino acids which are released in the stratum corneum contributing to water retention, ultraviolet B absorption and acidification.² Loss of function mutations in the gene encoding filaggrin (*FLG*) cause ichthyosis vulgaris³ and strongly predispose to eczema as a manifestation of skin barrier dysfunction.⁴ Whilst *FLG* null mutations lead to a marked reduction in filaggrin in skin, factors that increase the thickness of the stratum corneum may be considered to increase filaggrin expression in a specific or a non-specific way. Whilst filaggrin is intimately associated with cornification, it is interesting to note that *FLG* is still expressed in the non-keratinised areas of buccal mucosa.⁵

Lichen planus (LP) is an inflammatory disease of unknown aetiology. The increased expression of keratins 1, 2, 10 and 11, along with a reduction in expression of keratins 4 and 13 have been reported in buccal mucosal LP.⁶ It has been proposed that barrier impairment may be a preceding event in the pathogenesis of LP, or it may occur as a secondary effect resulting from disturbance in keratinocyte differentiation.⁷ Opposing hypotheses can therefore be proposed: that filaggrin expression is either increased or decreased within the lesional epidermis in LP.

The study by Larsen *et al.* reported in this edition of the *JEADV*⁸ set out to test several hypotheses relating to the relationship between oral lichenoid lesions (OLL, including oral LP), filaggrin expression, *FLG* mutations and co-associated skin conditions. 38 patients with OLL (including 19 with oral LP) and 11 with non-specific stomatitis were compared with 29 unaffected controls.

Immunofluorescent staining of buccal mucosal biopsy specimens showed a marked increase in filaggrin expression in 21% of patients with OLL but none of the healthy controls; conversely there was a marked reduction in filaggrin immunostaining observed in 15% of healthy controls compared with 6% of patients with OLL. The differences in filaggrin staining showed no association with the presence or absence of loss of function mutations in *FLG*. The patients with OLL also did not have an increased prevalence of dermatological disorders (other than LP) when compared with controls.

These findings must be interpreted with caution because of the small sample size. However, the observation of an increase in filaggrin expression in OLL that is independent of *FLG* genotype and unrelated to other skin disorders implies that the increased filaggrin expression is a secondary effect of local inflammation and hyperkeratosis rather than a primary aetiological factor. Filaggrin binds to keratins 2 and 10 and may therefore be increased in consort with them.⁶ The absence of association of LP with filaggrin down-regulation sheds doubt on the hypothesis that skin barrier dysfunction contributes to the primary pathogenesis of lichenoid inflammation; the underlying aetiology of oral LP continues to be enigmatic.

References

- 1 Sandilands A, Sutherland C, Irvine A *et al.* Filaggrin in the frontline: role in skin barrier function and disease. *J Cell Sci* 2009; **122**: 1285-94.
- 2 Brown SJ, McLean WH. One remarkable molecule: filaggrin. *J Invest Dermatol* 2012; **132**: 751-62.
- 3 Smith FJ, Irvine AD, Terron-Kwiatkowski A *et al.* Loss-of-function mutations in the gene encoding filaggrin cause ichthyosis vulgaris. *Nat Genet* 2006; **38**: 337-42.
- 4 Irvine AD, McLean WH, Leung DY. Filaggrin mutations associated with skin and allergic diseases. *N Engl J Med* 2011; **365**: 1315-27.
- 5 Smith SA, Dale BA. Immunologic localization of filaggrin in human oral epithelia and correlation with keratinization. *J Invest Dermatol* 1986; **86**: 168-72.
- 6 Boisnic S, Ouhayoun JP, Branchet MC *et al.* Alteration of cytokeratin expression in oral lichen planus. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 1995; **79**: 207-15.
- 7 Danielsson K, Ebrahimi M, Nylander E *et al.* Alterations in Factors Involved in Differentiation and Barrier Function in the Epithelium in Oral and Genital Lichen Planus. *Acta Derm Venereol* 2017; **97**: 214-8.
- 8 Larsen KR, Johansen JD, Reibel J *et al.* Filaggrin gene mutations and the distribution of filaggrin in oral mucosa of patients with oral lichen planus and healthy controls. *J Eur Acad Dermatol Venereol* 2016.